

Product datasheet for TA807495

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VEGFD Mouse Monoclonal Antibody [Clone ID: OTI3C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C3

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 89-205 of human

FIGF(NP_004460) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 37.8 kDa

Gene Name: c-fos induced growth factor

Database Link: NP 004460

Entrez Gene 2277 Human

O43915

VEGFD Mouse Monoclonal Antibody [Clone ID: OTI3C3] - TA807495

Background: The protein encoded by this gene is a member of the platelet-derived growth factor/vascular

endothelial growth factor (PDGF/VEGF) family and is active in angiogenesis,

lymphangiogenesis, and endothelial cell growth. This secreted protein undergoes a complex proteolytic maturation, generating multiple processed forms which bind and activate VEGFR-2 and VEGFR-3 receptors. This protein is structurally and functionally similar to vascular endothelial growth factor C. Read-through transcription has been observed between this

locus and the upstream PIR (GenelD 8544) locus. [provided by RefSeq, Feb 2011]


Synonyms: VEGF-D; VEGFD

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling

pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FIGF ([RC206549], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FIGF. Positive lysates [LY417969] (100ug) and [LC417969] (20ug) can be purchased separately from OriGene.