

### Product datasheet for TA807495AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **VEGFD Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3C3]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3C3

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse

**Isotype:** IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 89-205 of human

FIGF(NP\_004460) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 37.8 kDa

**Gene Name:** c-fos induced growth factor

Database Link: NP 004460

Entrez Gene 2277 Human

043915





**Background:** The protein encoded by this gene is a member of the platelet-derived growth factor/vascular

endothelial growth factor (PDGF/VEGF) family and is active in angiogenesis,

lymphangiogenesis, and endothelial cell growth. This secreted protein undergoes a complex proteolytic maturation, generating multiple processed forms which bind and activate VEGFR-2 and VEGFR-3 receptors. This protein is structurally and functionally similar to vascular endothelial growth factor C. Read-through transcription has been observed between this

locus and the upstream PIR (GenelD 8544) locus. [provided by RefSeq, Feb 2011]


Synonyms: VEGF-D; VEGFD

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Bladder cancer, Cytokine-cytokine receptor interaction, Focal adhesion, mTOR signaling

pathway, Pancreatic cancer, Pathways in cancer, Renal cell carcinoma

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FIGF ([RC206549], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FIGF. Positive lysates [LY417969] (100ug) and [LC417969] (20ug) can be purchased separately from OriGene.