

Product datasheet for TA807496BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DR4 (TNFRSF10A) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8A12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8A12

Applications: WB

Recommended Dilution: WB 1:200~2000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 263-468 of human

TNFRSF10A(NP_003835) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: tumor necrosis factor receptor superfamily member 10a

Database Link: NP 003835

Entrez Gene 8797 Human

O00220

Background: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice

suggested that FADD, a death domain containing adaptor protein, is required for the

apoptosis mediated by this protein. [provided by RefSeq, Jul 2008]

Synonyms: APO2; CD261; DR4; TRAILR-1; TRAILR1




DR4 (TNFRSF10A) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8A12] – TA807496BM


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TNFRSF10A (Cat# [RC202152], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF10A(Cat# [TA807496]).



Western blot analysis of extracts (10ug) from 1 cell line and 1 human tissue by using anti-TNFRSF10A monoclonal antibody at 1:200 dilution.