

## **Product datasheet for TA807511S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **EIF1AY Mouse Monoclonal Antibody [Clone ID: OTI2G2]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G2

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 2-144 of human

EIF1AY(NP\_004672) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 16.3 kDa

**Gene Name:** eukaryotic translation initiation factor 1A, Y-linked

Database Link: NP 004672

Entrez Gene 9086 Human

<u>014602</u>

**Background:** This gene is located on the non-recombining region of the Y chromosome. It encodes a

protein related to eukaryotic translation initiation factor 1A (EIF1A), which may function in stabilizing the binding of the initiator Met-tRNA to 40S ribosomal subunits. Alternative splicing


results in multiple transcript variants. [provided by RefSeq, Jul 2013]

Synonyms: eIF-4C






## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY EIF1AY (Cat# [RC202695], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-EIF1AY(Cat# [TA807511]). Positive lysates [LY417819] (100ug) and [LC417819] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 5 different cell lines by using anti-EIF1AY monoclonal antibody at 1:200 dilution.