

## **Product datasheet for TA807540S**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# TRIB3 Mouse Monoclonal Antibody [Clone ID: OTI2D11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D11

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human Host: Mouse

**Isotype:** IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-200 of human

TRIB3(NP\_066981) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39.4 kDa

**Gene Name:** tribbles pseudokinase 3

Database Link: NP 066981

Entrez Gene 57761 Human

Q96RU7

**Background:** The protein encoded by this gene is a putative protein kinase that is induced by the

transcription factor NF-kappaB. The encoded protein is a negative regulator of NF-kappaB and can also sensitize cells to TNF- and TRAIL-induced apoptosis. In addition, this protein can negatively regulate the cell survival serine-threonine kinase AKT1. Differential promoter usage and alternate splicing result in multiple transcript variants. [provided by RefSeq, Jul 2014]




### TRIB3 Mouse Monoclonal Antibody [Clone ID: OTI2D11] - TA807540S

**Synonyms:** C20orf97; NIPK; SINK; SKIP3; TRB3

**Protein Families:** Druggable Genome, Protein Kinase, Transcription Factors

# **Product images:**



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-TRIB3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA807540]) (1:150)