

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA807610AM

## RNF20 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2B9]

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI2B9
Applications:	IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 493-754 of human RNF20 (NP_062538) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	113.5 kDa
Gene Name:	ring finger protein 20
Database Link:	<u>NP_062538</u> <u>Entrez Gene 109331 MouseEntrez Gene 56254 Human</u> <u>Q5VTR2</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	RNF20 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2B9] – TA807610AM		
Background:	The protein encoded by this gene shares similarity with BRE1 of S. cerevisiae. The protein encoded by this human gene is an E3 ubiquitin ligase that regulates chromosome structure by monoubiquitinating histone H2B. This protein acts as a putative tumor suppressor and positively regulates the p53 tumor suppressor as well as numerous histone H2A and H2B genes. In contrast, this protein also suppresses the expression of several protooncogenes and growth-related genes, including many genes that are induced by epidermal growth factor. This gene selectively suppresses the expression of some genes by interfering with chromatin recruitment of transcription elongation factor SII (TFIIS). [provided by RefSeq, Feb 2012]		
Synonyms:	BRE1; BRE1A; hBRE1		


Protein Families:

Druggable Genome

### **Product images:**


170		
130	_	_
100		
70	-	
55		
40		
35		
25	—	
15	-1	
10	-1	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RNF20 (Cat# [RC218422], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RNF20 (Cat# [TA807610])(1:500). Positive lysates [LY412703] (100ug) and [LC412703] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-RNF20 mouse monoclonal antibody. (Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA807610]) (1:150)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-RNF20 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA807610]) (1:150)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US