

## Product datasheet for TA807610BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### RNF20 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2B9]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2B9
Applications: IHC, WB

**Reactivity:** WB 1:500, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 493-754 of human

RNF20 (NP\_062538) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 113.5 kDa

**Gene Name:** ring finger protein 20

Database Link: NP 062538

Entrez Gene 109331 MouseEntrez Gene 56254 Human

O5VTR2






### Background:

The protein encoded by this gene shares similarity with BRE1 of S. cerevisiae. The protein encoded by this human gene is an E3 ubiquitin ligase that regulates chromosome structure by monoubiquitinating histone H2B. This protein acts as a putative tumor suppressor and positively regulates the p53 tumor suppressor as well as numerous histone H2A and H2B genes. In contrast, this protein also suppresses the expression of several protooncogenes and growth-related genes, including many genes that are induced by epidermal growth factor. This gene selectively suppresses the expression of some genes by interfering with chromatin recruitment of transcription elongation factor SII (TFIIS). [provided by RefSeq, Feb 2012]


Synonyms: BRE1; BRE1A; hBRE1

Protein Families: Druggable Genome

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY RNF20 (Cat# [RC218422], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RNF20 (Cat# [TA807610])(1:500). Positive lysates [LY412703] (100ug) and [LC412703] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-RNF20 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA807610]) (1:150)





Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-RNF20 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA807610]) (1:150)