

# **Product datasheet for TA807650S**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## DR4 (TNFRSF10A) Mouse Monoclonal Antibody [Clone ID: OTI11B12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI11B12

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

**Isotype:** IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 263-468 of human

TNFRSF10A(NP\_003835) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Store** at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** tumor necrosis factor receptor superfamily member 10a

Database Link: NP 003835

Entrez Gene 8797 Human

O00220

**Background:** The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor

is activated by tumor necrosis factor-related apoptosis inducing ligand (TNFSF10/TRAIL), and thus transduces cell death signal and induces cell apoptosis. Studies with FADD-deficient mice

suggested that FADD, a death domain containing adaptor protein, is required for the

apoptosis mediated by this protein. [provided by RefSeq, Jul 2008]

**Synonyms:** APO2; CD261; DR4; TRAILR-1; TRAILR1






**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNFRSF10A ([RC202152], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNFRSF10A (1:2000).