

## **Product datasheet for TA807693**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

#### NDUFB11 Mouse Monoclonal Antibody [Clone ID: OTI8C5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI8C5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 30-163 of human

NDUFB11(NP\_061929) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 18.2 kDa

**Gene Name:** NADH:ubiquinone oxidoreductase subunit B11

Database Link: NP 061929

Entrez Gene 54539 Human

O9NX14

Background: NDUFB11 is a component of mitochondrial complex I. Complex I catalyzes the first step in the

electron transport chain, the transfer of 2 electrons from NADH to ubiquinone, coupled to the translocation of 4 protons across the membrane (Carroll et al., 2002 [PubMed 12381726]).

[supplied by OMIM, Feb 2009]


**Synonyms:** CI-ESSS; ESSS; Np15; NP17.3; P17.3






**Protein Families:** Transmembrane

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NDUFB11 ([RC213937], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NDUFB11. Positive lysates [LY412795] (100ug) and [LC412795] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 5 different cell lines and 4 human tissue by using anti-NDUFB11 monoclonal antibody (1: Hela; 2: HepG2; 3: K562; 4: Mcf7; 5: SVT2; 6: Liver; 7: Testis; 8: Brain; 9: Uterus) at 1:200 dilution.