

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA807759BM

### BAMBI Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D5]

### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI5D5
Applications:	WB
Recommended Dilution:	WB 1:200
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 21-152 of human BAMBI(NP_036474) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	26.6 kDa
Gene Name:	BMP and activin membrane-bound inhibitor
Database Link:	<u>NP_036474</u> <u>Entrez Gene 68010 MouseEntrez Gene 83837 RatEntrez Gene 25805 Human</u> <u>Q13145</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## SourcessBAMBI Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5D5] - TA807759BMBackground:This gene encodes a transmembrane glycoprotein related to the type I receptors of the<br/>transforming growth factor-beta (TGF-beta) family, whose members play important roles in<br/>signal transduction in many developmental and pathological processes. The encoded protein<br/>however is a pseudoreceptor, lacking an intracellular serine/threonine kinase domain<br/>required for signaling. Similar proteins in frog, mouse and zebrafish function as negative<br/>regulators of TGF-beta, which has led to the suggestion that the encoded protein may<br/>function to limit the signaling range of the TGF-beta family during early embryogenesis.<br/>[provided by RefSeq, Jul 2008]

Synonyms:	NMA
Protein Families:	Transmembrane

**Product images:** 

# 170 — 130 — 100 — 70 — 55 — 40 — 35 — 25 — 15 — 10 —

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY BAMBI (Cat# [RC200195], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-BAMBI (Cat# [TA807759])(1:2000). Positive lysates [LY415799] (100ug) and [LC415799] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US