

Product datasheet for TA807762M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PODXL Mouse Monoclonal Antibody [Clone ID: OTI9C2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9C2

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 32-461 of human

PODXL(NP_005388) produced in HEK293 cells.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: podocalyxin like

Database Link: NP 001018121

Entrez Gene 5420 Human

<u>O00592</u>



PODXL Mouse Monoclonal Antibody [Clone ID: OTI9C2] - TA807762M


Background:

This gene encodes a member of the sialomucin protein family. The encoded protein was originally identified as an important component of glomerular podocytes. Podocytes are highly differentiated epithelial cells with interdigitating foot processes covering the outer aspect of the glomerular basement membrane. Other biological activities of the encoded protein include: binding in a membrane protein complex with Na+/H+ exchanger regulatory factor to intracellular cytoskeletal elements, playing a role in hematopoetic cell differentiation, and being expressed in vascular endothelium cells and binding to L-selectin. [provided by RefSeq, Jul 2008]

Synonyms: gp135; Gp200; PC; PCLP; PCLP-1; PDX; PODXL1

Protein Families: Transmembrane

Product images:



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-PODXL mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.