

## Product datasheet for TA807790BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Myostatin Propeptide (MSTN) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI6A7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6A7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 24-266 of human

MSTN(NP\_005250) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 42.6 kDa

Gene Name: myostatin

Database Link: NP 005250

Entrez Gene 17700 MouseEntrez Gene 29152 RatEntrez Gene 2660 Human

014793





# Myostatin Propeptide (MSTN) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI6A7] – TA807790BM

**Background:** The protein encoded by this gene is a member of the bone morphogenetic protein (BMP)


family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. The members of this family are regulators of cell growth and differentiation in both embryonic and adult tissues. This gene is thought to encode a secreted protein which negatively regulates skeletal muscle growth. [provided by RefSeq, Jul

2008]

Synonyms: GDF8; MSLHP

**Protein Families:** Druggable Genome, Secreted Protein

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MSTN ([RC210368], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MSTN (1:2000). Positive lysates [LY401622] (100ug) and [LC401622] (20ug) can be purchased separately from OriGene.