

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA807821

TNF alpha (TNF) Mouse Monoclonal Antibody [Clone ID: OTI5A11]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI5A11
Applications:	IF, WB
Recommended Dilution:	WB 1:2000, IF 1:100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TNFA (NP_000585) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25.5 kDa
Gene Name:	tumor necrosis factor
Database Link:	<u>NP_000585</u> <u>Entrez Gene 7124 Human</u> <u>P01375</u>
Synonyms:	DIF; TNF-alpha; TNFA; TNFSF2
Protein Families:	Druggable Genome, Secreted Protein, Transcription Factors, Transmembrane




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


STANDARIGENE TNF alpha (TNF) Mouse Monoclonal Antibody [Clone ID: OTI5A11] – TA807821 TA807821

Protein Pathways:Adipocytokine signaling pathway, Allograft rejection, Alzheimer's disease, Amyotrophic lateral
sclerosis (ALS), Apoptosis, Asthma, Cytokine-cytokine receptor interaction, Dilated
cardiomyopathy, Fc epsilon RI signaling pathway, Graft-versus-host disease, Hematopoietic
cell lineage, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway, Natural killer cell
mediated cytotoxicity, NOD-like receptor signaling pathway, RIG-I-like receptor signaling
pathway, Systemic lupus erythematosus, T cell receptor signaling pathway, TGF-beta signaling
pathway, Toll-like receptor signaling pathway, Type I diabetes mellitus, Type II diabetes
mellitus

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (middle lane) or pCMV6-ENTRY TNFA ([RC206983], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) and Origene recombinant TNFA protein [TP750007] (0.6ug, left lane) were separated by SDS-PAGE and immunoblotted with anti-TNFA (1:2000). Positive lysates [LY424626] (100ug) and [LC424626] (20ug) can be purchased separately from OriGene.



ELISA anaylysis of anti-TNFA antibody. 96-well plate was coated with Origene recombinant protein TNFa ([TP750007]) and primary antibody dilution ranges from 1:100 to 1:10000 (TA807821, 1mg/ml).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US