

Product datasheet for TA807821M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TNF alpha (TNF) Mouse Monoclonal Antibody [Clone ID: OTI5A11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5A11
Applications: IF, WB

Recommended Dilution: WB 1:2000, IF 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TNFA (NP_000585) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 25.5 kDa

Gene Name: tumor necrosis factor

Database Link: NP 000585

Entrez Gene 7124 Human

P01375

Synonyms: DIF; TNF-alpha; TNFA; TNFSF2


Protein Families: Druggable Genome, Secreted Protein, Transcription Factors, Transmembrane




Protein Pathways:

Adipocytokine signaling pathway, Allograft rejection, Alzheimer's disease, Amyotrophic lateral sclerosis (ALS), Apoptosis, Asthma, Cytokine-cytokine receptor interaction, Dilated cardiomyopathy, Fc epsilon RI signaling pathway, Graft-versus-host disease, Hematopoietic cell lineage, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway, Natural killer cell mediated cytotoxicity, NOD-like receptor signaling pathway, RIG-I-like receptor signaling pathway, Systemic lupus erythematosus, T cell receptor signaling pathway, TGF-beta signaling pathway, Toll-like receptor signaling pathway, Type I diabetes mellitus, Type II diabetes mellitus

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (middle lane) or pCMV6-ENTRY TNFA ([RC206983], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) and Origene recombinant TNFA protein [TP750007] (0.6ug, left lane) were separated by SDS-PAGE and immunoblotted with anti-TNFA (1:2000). Positive lysates [LY424626] (100ug) and [LC424626] (20ug) can be purchased separately from OriGene.



ELISA analysis of anti-TNFA antibody. 96-well plate was coated with Origene recombinant protein TNFa ([TP750007]) and primary antibody dilution ranges from 1:100 to 1:10000 ([TA807821], 1mg/ml).