

Product datasheet for TA807959

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cardiac Troponin T (TNNT2) Mouse Monoclonal Antibody [Clone ID: OTI8B5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8B5
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Native troponin complex

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: troponin T2, cardiac type

Database Link: NP 000355

Entrez Gene 21956 MouseEntrez Gene 24837 RatEntrez Gene 7139 Human

P45379

Background: Troponin I (TnI), along with troponin T (TnT) and troponin C (TnC), is one of 3 subunits that

form the troponin complex of the thin filaments of striated muscle. Tnl is the inhibitory subunit; blocking actin-myosin interactions and thereby mediating striated muscle relaxation. The Tnl subfamily contains three genes: Tnl-skeletal-fast-twitch, Tnl-skeletal-slow-twitch, and Tnl-cardiac. This gene encodes the Tnl-cardiac protein and is exclusively expressed in cardiac muscle tissues. Mutations in this gene cause familial hypertrophic cardiomyopathy type 7

(CMH7) and familial restrictive cardiomyopathy (RCM). [provided by RefSeq, Jul

Synonyms: CMD1D; CMH2; CMPD2; cTnT; LVNC6; RCM3; TnTC







Protein Families: Druggable Genome

Protein Pathways: Cardiac muscle contraction, Dilated cardiomyopathy, Hypertrophic cardiomyopathy (HCM)

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TNNT2 ([RC214180], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TNNT2. Positive lysates [LY400130] (100ug) and [LC400130] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human adult heart tissue using anti-TNNT2 mouse monoclonal antibody. (Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA807959)