

# Product datasheet for TA808069AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### PD1 (PDCD1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI22A9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI22A9

Applications: FC, IF, WB

**Recommended Dilution:** WB 1:500, IF 1:100~900, FLOW 1:50

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDCD1 (NP\_005009) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 29.2 kDa

**Gene Name:** programmed cell death 1

Database Link: NP 005009

Entrez Gene 5133 Human

015116





**Background:** This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This


protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul 2008]

Synonyms: CD279; hPD-1; hPD-1; hSLE1; PD-1; PD1; SLEB2


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs), T cell receptor signaling pathway


## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1 ([RC210364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1 (1:500). Positive lysates [LY401555] (100ug) and [LC401555] (20ug) can be purchased separately from OriGene.




Immunofluorescent staining of PDCD1 ([RC210364])-stable-transfected HEK293T cells (left) labeling PDCD1 with mouse monoclonal antibody [TA808069] (green) and nucleus with Hoechst33342 (blue). HEK293T cells serve as negative control (right) (1:100).




293T cell line



Immunocytochemistry staining of stable expression PD1 cells using anti-PDCD1 mouse monoclonal antibody ([TA808069]) (Left). The rihgt is negative control (1:900).



Flow cytometric Analysis of stable expression PD1 cells using anti-PDCD1 antibody ([TA808069]) (Red) compared to a nonspecific negative control antibody (Blue) (1:50).



Flow cytometric Analysis of stable expression PDL1 ([RC213071]) cells using anti-PDCD1 antibody ([TA808069]) (blue) or 0.3ug/ml PD1-Fc fusion protein ([TP700199]) (green) or both (red), and detected by anti-Fc (human) IgG-FITC (1:50).