

Product datasheet for TA808081

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL12A Mouse Monoclonal Antibody [Clone ID: OTI2A8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2A8

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 57-253 of human

IL12A(NP_000873) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: interleukin 12A

Database Link: NP 000873

Entrez Gene 3592 Human

P29459



Background:

This gene encodes a subunit of a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. The cytokine is a disulfide-linked heterodimer composed of the 35-kD subunit encoded by this gene, and a 40-kD subunit that is a member of the cytokine receptor family. This cytokine is required for the T-cell-independent induction of interferon (IFN)-gamma, and is important for the differentiation of both Th1 and Th2 cells. The responses of lymphocytes to this cytokine are mediated by the activator of transcription protein STAT4. Nitric oxide synthase 2A (NOS2A/NOS2) is found to be required for the signaling process of this cytokine in innate immunity. [provided by RefSeq, Jul 2008]

Synonyms: CLMF; IL-12A; NFSK; NKSF1; P35

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: Allograft rejection, Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway, RIG-I-

like receptor signaling pathway, Toll-like receptor signaling pathway, Type I diabetes mellitus

Product images:




Figure A, Western blot analysis of overexpressed lysates(25ug per lane) from HEK293T cells transfected with empty plasmid ([PS100001], lane 1), human IL12A plasmid ([RC211224], lane 2), mouse IL12A plasmid ([MR225609], lane 3), rat IL12A plasmid ([RR210914], lane 4) using anti-IL12A antibody TA808081 (1:500). Figure B, Western blot analysis of the same samples as figure A with anti-DDK antibody ([TA180144], 1:1000)