

# **Product datasheet for TA808195M**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **IL12RB1 Mouse Monoclonal Antibody [Clone ID: OTI1A6]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1A6

Applications: WB

Reactivity: WB 1:500

Human

Host: Mouse

Isotype: IgG1
Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 33-283 of human

IL12RB1(NP 714912) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39.6 kDa

**Gene Name:** interleukin 12 receptor subunit beta 1

Database Link: NP 714912

Entrez Gene 3594 Human

P42701



Background:


The protein encoded by this gene is a type I transmembrane protein that belongs to the hemopoietin receptor superfamily. This protein binds to interleukine 12 (IL12) with a low affinity, and is thought to be a part of IL12 receptor complex. This protein forms a disulfide-linked oligomer, which is required for its IL12 binding activity. The coexpression of this and IL12RB2 proteins was shown to lead to the formation of high-affinity IL12 binding sites and reconstitution of IL12 dependent signaling. Mutations in this gene impair the development of interleukin-17-producing T lymphocytes and result in increased susceptibility to mycobacterial and Salmonella infections. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2014]

Synonyms: CD212; IL-12R-BETA1; IL12RB; IMD30

Protein Families: Druggable Genome, Transmembrane

**Protein Pathways:** Cytokine-cytokine receptor interaction, Jak-STAT signaling pathway

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IL12RB1 (Cat# [RC216444], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IL12RB1 (Cat# [TA808195])(1:500). Positive lysates [LY403516] (100ug) and [LC403516] (20ug) can be purchased separately from OriGene.