

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TA808223S

# CDC42 binding protein kinase alpha (CDC42BPA) Mouse Monoclonal Antibody [Clone ID: OTI7B9]

### **Product data:**

Product Type:	Primary Antibodies	
Clone Name:	OTI7B9	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1419-1638 of human CDC42BPA(NP_055641) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	185.9 kDa	
Gene Name:	CDC42 binding protein kinase alpha	
Database Link:	<u>NP_055641</u> <u>Entrez Gene 114116 RatEntrez Gene 226751 MouseEntrez Gene 8476 Human</u> <u>Q5VT25</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	CDC42 binding protein kinase alpha (CDC42BPA) Mouse Monoclonal Antibody [Clone ID: OTI7B9] – TA808223S
Background:	The protein encoded by this gene is a member of the Serine/Threonine protein kinase family. This kinase contains multiple functional domains. Its kinase domain is highly similar to that of the myotonic dystrophy protein kinase (DMPK). This kinase also contains a Rac interactive binding (CRIB) domain, and has been shown to bind CDC42. It may function as a CDC42 downstream effector mediating CDC42 induced peripheral actin formation, and promoting cytoskeletal reorganization. Multiple alternatively spliced transcript variants have been described, and the full-length nature of two of them has been reported. [provided by RefSeq, Jul 2008]
Synonyms:	MRCK; MRCKA; PK428
Protein Familie	: Druggable Genome, Protein Kinase

### **Product images:**

170	-	
130	_	
100		
70	-	
55	-	
40		
35	-	
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CDC42BPA (Cat# [RC222587], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CDC42BPA (Cat# [TA808223])(1:2000). Positive lysates [LY415030] (100ug) and [LC415030] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US