

## Product datasheet for TA808230BM

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PRPK (TP53RK) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5G7]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5G7
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human TP53RK(NP\_291028) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 28 kDa

**Gene Name:** TP53 regulating kinase

Database Link: NP 291028

Entrez Gene 112858 Human

Q96S44


Synonyms: BUD32; C20orf64; dJ101A2; Nori-2; Nori-2p; PRPK

**Protein Families:** Druggable Genome, Protein Kinase






## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TP53RK ([RC209748], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TP53RK (1:2000). Positive lysates [LY403262] (100ug) and [LC403262] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-TP53RK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA808230]) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-TP53RK mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA808230]) (1:150)