

Product datasheet for TA808235BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ZNF69 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI3D11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3D11
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ZNF69(NP_068734) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 17.2 kDa

Gene Name: zinc finger protein 69

Database Link: NP 068734

Entrez Gene 7620 Human

Q9UC07

Background: May be involved in transcriptional regulation. [UniProtKB/Swiss-Prot Function]


Synonyms: Cos5; hZNF3

Protein Families: Transcription Factors






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ZNF69 ([RC208624], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ZNF69 (1:2000). Positive lysates [LY411885] (100ug) and [LC411885] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-ZNF69 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA808235]) (1:150)





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-ZNF69 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA808235]) (1:150)