

Product datasheet for TA808436M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SCMH1 Mouse Monoclonal Antibody [Clone ID: OTI7B6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7B6
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-309 of human

SCMH1(NP_036368) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 65.9 kDa

Gene Name: sex comb on midleg homolog 1 (Drosophila)

Database Link: NP 036368

Entrez Gene 22955 Human

Q96GD3

Background: Interacts with the SAM domain of PHC1 via its SAM domain in vitro (By similarity). Associates

with a PRC1-like complex. [UniProtKB/Swiss-Prot Function]


Synonyms: Scml3

Protein Families: Transcription Factors







Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SCMH1 ([RC200193], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SCMH1 (1:500). Positive lysates [LY415889] (100ug) and [LC415889] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-SCMH1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-SCMH1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.