

Product datasheet for TA808449BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

NOBOX Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8C7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8C7
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-271 of human NOBOX

(NP_001073882) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 61.1 kDa

Gene Name: NOBOX oogenesis homeobox

Database Link: NP 001073882

Entrez Gene 135935 Human

060393

Background: This homeobox gene encodes a transcription factor that is thought to play a role in

oogenesis. In mice, it is essential for folliculogenesis and regulation of oocyte-specific genes. Defects in this gene result in premature ovarian failure type 5. [provided by RefSeq, May


2011]

Synonyms: OG-2; OG2; OG2X; POF5; TCAG 12042






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NOBOX ([RC216621], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NOBOX (1:2000). Positive lysates [LY421630] (100ug) and [LC421630] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-NOBOX mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.