

Product datasheet for TA808481BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TRBP (TARBP2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI7C1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI7C1
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TARBP2 (NP_004169) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36.7 kDa

Gene Name: TARBP2, RISC loading complex RNA binding subunit

Database Link: NP 004169

Entrez Gene 21357 MouseEntrez Gene 363006 RatEntrez Gene 6895 Human

Q15633






Background:

HIV-1, the causative agent of acquired immunodeficiency syndrome (AIDS), contains an RNA genome that produces a chromosomally integrated DNA during the replicative cycle. Activation of HIV-1 gene expression by the transactivator Tat is dependent on an RNA regulatory element (TAR) located downstream of the transcription initiation site. The protein encoded by this gene binds between the bulge and the loop of the HIV-1 TAR RNA regulatory element and activates HIV-1 gene expression in synergy with the viral Tat protein. Alternative splicing results in multiple transcript variants encoding different isoforms. This gene also has a pseudogene. [provided by RefSeq, Jul 2008]


Synonyms: LOQS; TRBP; TRBP1; TRBP2

Protein Families: Stem cell - Pluripotency

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TARBP2 (Cat# [RC201043], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TARBP2 (Cat# [TA808481])(1:2000). Positive lysates [LY418167] (100ug) and [LC418167] (20ug) can be purchased separately from OriGene.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and TARBP2-Knockout HeLa cells (KO, Cat# [LC833009]) were separated by SDS-PAGE and immunoblotted with anti-TARBP2 monoclonal antibody [TA808481] (1:500`). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-TARBP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA, pH8.0, [TA808481]) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-TARBP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA, pH8.0, [TA808481]) (1:150)



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-TARBP2 mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA, pH8.0, [TA808481]) (1:150)