

# **Product datasheet for TA808493**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **AMACR Mouse Monoclonal Antibody [Clone ID: OTI6C7]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6C7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

**Isotype:** lgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human AMACR (NP\_055139) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 42.2 kDa

**Gene Name:** alpha-methylacyl-CoA racemase

Database Link: NP 055139

Entrez Gene 23600 Human

O9UHK6





Background:


This gene encodes a racemase. The encoded enzyme interconverts pristanoyl-CoA and C27-bile acylCoAs between their (R)- and (S)-stereoisomers. The conversion to the (S)-stereoisomers is necessary for degradation of these substrates by peroxisomal beta-oxidation. Encoded proteins from this locus localize to both mitochondria and peroxisomes. Mutations in this gene may be associated with adult-onset sensorimotor neuropathy, pigmentary retinopathy, and adrenomyeloneuropathy due to defects in bile acid synthesis. Alternatively spliced transcript variants have been described. Read-through transcription also exists between this gene and the upstream neighboring C1QTNF3 (C1q and tumor necrosis factor related protein 3) gene. [provided by RefSeq, Mar 2011]

Synonyms: AMACRD; CBAS4; RACE; RM

**Protein Families:** Druggable Genome

**Protein Pathways:** Metabolic pathways, Primary bile acid biosynthesis

### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY AMACR ([RC213437], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-AMACR (1:2000). Positive lysates [LY402314] (100ug) and [LC402314] (20ug) can be purchased separately from OriGene.