

Product datasheet for TA808562AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ERCC8 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI9B11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9B11

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ERCC8(NP_001007235) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 43.9 kDa

Gene Name: ERCC excision repair 8, CSA ubiquitin ligase complex subunit

Database Link: NP 001007235

Entrez Gene 71991 MouseEntrez Gene 310071 RatEntrez Gene 1161 Human

Q13216

Background: This gene encodes a WD repeat protein, which interacts with Cockayne syndrome type B

(CSB) protein and with p44 protein, a subunit of the RNA polymerase II transcription factor IIH. Mutations in this gene have been identified in patients with hereditary disease Cockayne syndrome (CS). CS cells are abnormally sensitive to ultraviolet radiation and are defective in the repair of transcriptionally active genes. Several transcript variants encoding different

isoforms have been found for this gene. [provided by RefSeq, Mar 2014]






Synonyms: CKN1; CSA; UVSS2

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Nucleotide excision repair, Ubiquitin mediated proteolysis

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ERCC8 ([RC203124], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ERCC8 (1:2000). Positive lysates [LY423459] (100ug) and [LC423459] (20ug) can be purchased separately from OriGene.