

Product datasheet for **TA808610AM**

Thymine DNA glycosylase (TDG) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3C11]

Product data:

| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | OTI3C11 |
| Applications: | IHC, WB |
| Recommended Dilution: | WB 1:2000, IHC 1:150 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Human recombinant protein fragment corresponding to amino acids 160-253 of human TDG(NP_003202) produced in E.coli. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 0.5 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Biotin |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 45.9 kDa |
| Gene Name: | thymine DNA glycosylase |
| Database Link: | NP_003202 Entrez Gene 21665 Mouse Entrez Gene 114521 Rat Entrez Gene 6996 Human Q13569 |

[View online »](#)

Background:

The protein encoded by this gene belongs to the TDG/mug DNA glycosylase family. Thymine-DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired thymine. With lower activity, this enzyme also removes thymine from C/T and T/T mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. This enzyme plays a central role in cellular defense against genetic mutation caused by the spontaneous deamination of 5-methylcytosine and cytosine. This gene may have a pseudogene in the p arm of chromosome 12. [provided by RefSeq, Jul 2008]

Synonyms:

hTDG


Protein Families:

Druggable Genome


Protein Pathways:

Base excision repair


Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TDG (Cat# [RC207113], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TDG (Cat# [TA808610])(1:2000). Positive lysates [LY401109] (100ug) and [LC401109] (20ug) can be purchased separately from OriGene.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and TDG-Knockout HeLa cells (KO, Cat# [LC810235]) were separated by SDS-PAGE and immunoblotted with anti-TDG monoclonal antibody [TA808610], (1:500). Then the blotted membrane was stripped and reprobed with anti-beta-actin antibody ([TA811000]) as a loading control.




Immunohistochemical staining of paraffin-embedded Human colon tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human kidney tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human pancreas tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human thyroid tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human prostate tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Carcinoma of Human prostate tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffin-embedded Human bladder tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-TDG mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.