

Product datasheet for TA808784

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

MAK Mouse Monoclonal Antibody [Clone ID: OTI4H4]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: OTI4H4

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Rat

Host: Mouse

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 291-623 of human

MAK(NP 005897) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: male germ cell associated kinase

lgG1

Database Link: NP 005897

Entrez Gene 25677 RatEntrez Gene 4117 Human

P20794






Background:

The product of this gene is a serine/threonine protein kinase related to kinases involved in cell cycle regulation. It is expressed almost exclusively in the testis, primarily in germ cells. Studies of the mouse and rat homologs have localized the kinase to the chromosomes during meiosis in spermatogenesis, specifically to the synaptonemal complex that exists while homologous chromosomes are paired. There is, however, a study of the mouse homolog that has identified high levels of expression in developing sensory epithelia so its function may be more generalized. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2011]

Synonyms: dJ417M14.2; RP62

Protein Families: Druggable Genome, Protein Kinase

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAK ([RC213532], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAK (1:500).