

Product datasheet for TA808810S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SHMT2 Mouse Monoclonal Antibody [Clone ID: OTI3B3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3B3
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SHMT2(NP_005403) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 52.5 kDa

Gene Name: serine hydroxymethyltransferase 2

Database Link: NP 005403

Entrez Gene 108037 MouseEntrez Gene 299857 RatEntrez Gene 6472 Human

P34897

Background: This gene encodes the mitochondrial form of a pyridoxal phosphate-dependent enzyme that

catalyzes the reversible reaction of serine and tetrahydrofolate to glycine and 5,10-methylene tetrahydrofolate. The encoded product is primarily responsible for glycine synthesis. The activity of the encoded protein has been suggested to be the primary source of intracellular glycine. The gene which encodes the cytosolic form of this enzyme is located on chromosome 17. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009]




Synonyms: GLYA; HEL-S-51e; NEDCASB; SHMT

Protein Pathways: Cyanoamino acid metabolism, Glycine, serine and threonine metabolism, Metabolic

pathways, Methane metabolism, One carbon pool by folate

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SHMT2 ([RC204239], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SHMT2 (1:2000). Positive lysates [LY401661] (100ug) and [LC401661] (20ug) can be purchased separately from OriGene.