

### **Product datasheet for TA808811S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Inositol Hexakisphosphate Kinase 2 (IP6K2) Mouse Monoclonal Antibody [Clone ID: OTI1F12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1F12

Applications: WB

Recommended Dilution: WB 1:500

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1-250 of human

IP6K2(NP\_057375) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** inositol hexakisphosphate kinase 2

Database Link: NP 057375

Entrez Gene 59268 RatEntrez Gene 76500 MouseEntrez Gene 51447 Human

Q9UHH9

**Background:** This gene encodes a protein that belongs to the inositol phosphokinase (IPK) family. This

protein is likely responsible for the conversion of inositol hexakisphosphate (InsP6) to diphosphoinositol pentakisphosphate (InsP7/PP-InsP5). It may also convert 1,3,4,5,6-pentakisphosphate (InsP5) to PP-InsP4 and affect the growth suppressive and apoptotic activities of interferon-beta in some ovarian cancers. Alternative splicing results in multiple

transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008]

Synonyms: IHPK2; PIUS






**Protein Families:** 

Druggable Genome

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IP6K2 (Cat# [RC209533], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IP6K2 (1:500) (Cat# [TA808811]). Positive lysates [LY414064] (100ug) and [LC414064] (20ug) can be purchased separately from OriGene.