

## Product datasheet for **TA808934**


### TRIM56 Mouse Monoclonal Antibody [Clone ID: OTI8E12]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	OTI8E12
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 311-554 of human TRIM56(NP_112223) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	81.3 kDa
Gene Name:	tripartite motif containing 56
Database Link:	<a href="#">NP_112223</a> <a href="#">Entrez Gene 81844 Human</a> <a href="#">Q9BRZ2</a>
Background:	E3 ubiquitin-protein ligase that mediates 'Lys-63'-linked polyubiquitination of TMEM173/STING, thereby playing a key role in innate immunity. TMEM173/STING 'Lys-63'-linked ubiquitination activates the production of type I interferon IFN-beta following detection of pathogen- and host-derived double-stranded DNA. [UniProtKB/Swiss-Prot Function]
Synonyms:	RNF109




[View online »](#)

**Product images:**

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TRIM56 ([RC200891], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TRIM56 (1:2000). Positive lysates [LY410625] (100ug) and [LC410625] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-TRIM56 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808934) (1:150)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-TRIM56 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA808934) (1:150)