

Product datasheet for TA809142S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GATA3 Mouse Monoclonal Antibody [Clone ID: OTI8E11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8E11
Applications: FC, IF, WB
Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 155-443 of human

GATA3 (NP_001002295) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 47.9 kDa

Gene Name: GATA binding protein 3

Database Link: NP 001002295

Entrez Gene 14462 MouseEntrez Gene 85471 RatEntrez Gene 2625 Human

P23771

Background: This gene encodes a protein which belongs to the GATA family of transcription factors. The

protein contains two GATA-type zinc fingers and is an important regulator of T-cell

development and plays an important role in endothelial cell biology. Defects in this gene are the cause of hypoparathyroidism with sensorineural deafness and renal dysplasia. [provided

by RefSeq, Nov 2009]





Synonyms:


HDR; HDRS


Protein Families:

Adult stem cells, ES Cell Differentiation/IPS, Stem cell relevant signaling - JAK/STAT signaling

pathway, Transcription Factors

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GATA3 ([RC211904], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GATA3 (1:2000). Positive lysates [LY424146] (100ug) and [LC424146] (20ug) can be purchased separately from OriGene.

Western blot analysis of extracts (35ug) from 3 different cell lines by using anti-GATA3 monoclonal antibody (1:200).



MCF7

HUVEC






Immunocytochemistry staining of MCF-7 cells using anti-GATA3 mouse monoclonal antibody ([TA809142]) (Left). The right is negative control (1:1000).


MDA-MB-468


HUVEC





Immunocytochemistry staining of MDA-MB-468 cells using anti-GATA3 mouse monoclonal antibody ([TA809142]) (Left). The right is negative control (1:1000).





Flow cytometric Analysis of penetrated MCF-7 cells, using anti-GATA3 antibody ([TA809142]), (Red), compared to a nonspecific negative control antibody, (Blue). The right is HUVEC cells as negtive control (1:100).