

## **Product datasheet for TA809177**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Alkaline Phosphatase (ALPI) Mouse Monoclonal Antibody [Clone ID: OTI3D5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3D5
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human ALPI (NP\_001622) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 54.7 kDa

**Gene Name:** alkaline phosphatase, intestinal

Database Link: NP 001622

Entrez Gene 248 Human

P09923

**Background:** There are at least four distinct but related alkaline phosphatases: intestinal, placental,

placental-like, and liver/bone/kidney (tissue non-specific). The intestinal alkaline phosphatase gene encodes a digestive brush-border enzyme. This enzyme is upregulated during small

intestinal epithelial cell differentiation. [provided by RefSeq, Jul 2008]

Synonyms: IAP







**Protein Families:** Druggable Genome

**Protein Pathways:** Folate biosynthesis, Metabolic pathways

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ALPI ([RC224178], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ALPI (1:2000). Positive lysates [LY400612] (100ug) and [LC400612] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-ALPI mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.