

# Product datasheet for TA809201M

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Tau (MAPT) Mouse Monoclonal Antibody [Clone ID: OTI13B5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI13B5
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:500 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 623-758 of human

MAPT(NP\_058519) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 78.7 kDa

**Gene Name:** microtubule associated protein tau

Database Link: NP 058519

Entrez Gene 17762 MouseEntrez Gene 29477 RatEntrez Gene 4137 Human

P10636



## Tau (MAPT) Mouse Monoclonal Antibody [Clone ID: OTI13B5] - TA809201M

Background: This gene encodes the microtubule-associated protein tau (MAPT) whose transcript

undergoes complex, regulated alternative splicing, giving rise to several mRNA species. MAPT transcripts are differentially expressed in the nervous system, depending on stage of neuronal maturation and neuron type. MAPT gene mutations have been associated with several neurodegenerative disorders such as Alzheimer's disease, Pick's disease,

frontotemporal dementia, cortico-basal degeneration and progressive supranuclear palsy.


[provided by RefSeq, Jul 2008]

Synonyms: DDPAC; FTDP-17; MAPTL; MSTD; MTBT1; MTBT2; PPND; PPP1R103; TAU; tau-40


**Protein Families:** Druggable Genome

**Protein Pathways:** Alzheimer's disease, MAPK signaling pathway

# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY MAPT (Cat# [RC216166], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAPT (Cat# [TA809201])(1:2000). Positive lysates [LY429527] (100ug) and [LC429527] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 2 different cell lines by using anti-MAPT monoclonal antibody (1:500).





Western blot analysis of extracts (35ug) from human brain tissue by using anti-MAPT monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded Human embryonic cerebellum within the normal limits using anti-MAPT mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.