

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA809306M

### Bile Acid Receptor (NR1H4) Mouse Monoclonal Antibody [Clone ID: OTI4F12]

#### **Product data:**

Product Type:	Primary Antibodies	
Clone Name:	OTI4F12	
Applications:	IHC, WB	
Recommended Dilution:	WB 1:2000, IHC 1:150	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 21-297 of human NR1H4 (NP_005114) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Gene Name:	nuclear receptor subfamily 1 group H member 4	
Database Link:	<u>NP_005114</u> Entrez Gene 20186 MouseEntrez Gene 60351 RatEntrez Gene 9971 Human Q96RI1	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Bile Acid Receptor (NR1H4) Mouse Monoclonal Antibody [Clone ID: OTI4F12] – TA809306M	
Background:	This gene encodes a ligand-activated transcription factor, which shares structural features in common with nuclear hormone receptor family, such as a DNA-binding domain that targets the receptor to specific DNA sequences, and a ligand-binding domain, which interacts directly with the ligand and contains a ligand-dependent transcriptional activation domain. This protein functions as a receptor for bile acids, and when bound to bile acids, regulates the expression of genes involved in bile acid synthesis and transport. Alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Aug 2011]	
Synonyms:	BAR; FXR; HRR-1; HRR1; PFIC5; RIP14	
<b>Protein Families</b>	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors	

## **Product images:**

- 🕅 -

170	_	
130	-	
100		
70	-	
55		
40		
35		
25	-	
15	-	
10	-	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NR1H4 (Cat# [RC217443], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NR1H4 (Cat# [TA809306])(1:2000). Positive lysates [LY401577] (100ug) and [LC401577] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-NR1H4 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US