

## Product datasheet for **TA809308BM**

### **HNRNPA0 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8H8]**

#### **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI8H8
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 139-183 of human HNRNPA0 (NP_006796) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	30.7 kDa
Gene Name:	heterogeneous nuclear ribonucleoprotein A0
Database Link:	<a href="#">NP_006796</a> <a href="#">Entrez Gene 10949 Human</a> <a href="#">Q13151</a>




[View online »](#)

**Background:**


This gene belongs to the A/B subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene has two repeats of quasi-RRM domains that bind RNAs, followed by a glycine-rich C-terminus. [provided by RefSeq, Jul 2008]

**Synonyms:**


HNRPA0

**Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HNRNPA0 (Cat# [RC203666], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HNRNPA0 (Cat# [TA809308])(1:2000). Positive lysates [LY416409] (100ug) and [LC416409] (20ug) can be purchased separately from OriGene.




Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and HNRNPA0-Knockout 293T cells (KO, Cat# [LC811683]) were separated by SDS-PAGE and immunoblotted with anti-HNRNPA0 monoclonal antibody [TA809308], (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90AB1 antibody ([TA500494]) as a loading control.




Immunohistochemical staining of paraffin-embedded Human breast tissue within the normal limits using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human thyroid tissue using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)




Immunohistochemical staining of paraffin-embedded Human endometrium tissue within the normal limits using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human endometrium tissue using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)



Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)



Immunohistochemical staining of paraffin-embedded Human lymphoma tissue using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)



Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-HNRNPA0 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809308]) (1:150)