

## **Product datasheet for TA809334S**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# IRF3 Mouse Monoclonal Antibody [Clone ID: OTI2G6]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2G6
Applications: IF, WB
Recommended Dilution: WB 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 27-169 of human

IRF3(NP\_001184053) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** interferon regulatory factor 3

Database Link: NP 001184053

Entrez Gene 3661 Human

Q14653

**Background:** This gene encodes a member of the interferon regulatory transcription factor (IRF) family.

The encoded protein is found in an inactive cytoplasmic form that upon serine/threonine phosphorylation forms a complex with CREBBP. This complex translocates to the nucleus and activates the transcription of interferons alpha and beta, as well as other interferon-induced

 $genes.\ Alternatively\ spliced\ transcript\ variants\ encoding\ multiple\ isoforms\ have\ been$ 

observed for this gene. [provided by RefSeq, Nov 2011]

Synonyms: IIAE7






**Protein Families:** Druggable Genome, Transcription Factors


Protein Pathways: Cytosolic DNA-sensing pathway, RIG-I-like receptor signaling pathway, Toll-like receptor

signaling pathway

### **Product images:**





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IRF3 ([RC209951], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IRF3 (1:500).




Western blot analysis of extracts (35ug) from 6 different cell lines and 2 tissue lysates by using anti-IRF3 monoclonal antibody (1:500).









Immunofluorescent staining of 293T cells transfected by pCMV6-ENTRY IRF3 ([RC209951]) using anti-IRF3 antibody ([TA809334]/green, upper left; DAPI/blue, lower left; MERGED, upper right). 293T cells transfected with empty vector served as a negative control (MERGED, lower right) (1:100).