

## Product datasheet for **TA809448M**

### DIRAS2 Mouse Monoclonal Antibody [Clone ID: OTI10G5]


#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI10G5
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-196 of human DIRAS2(NP_060064) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	22.3 kDa
Gene Name:	DIRAS family GTPase 2
Database Link:	<a href="#">NP_060064</a> <a href="#">Entrez Gene 54769 Human</a> <a href="#">Q96HU8</a>
Background:	DIRAS2 belongs to a distinct branch of the functionally diverse Ras (see HRAS; MIM 190020) superfamily of monomeric GTPases. [supplied by OMIM, Apr 2004]. ##Evidence-Data-START## Transcript exon combination :: AK122986.1, BC008065.2 [ECO:0000332] RNAseq introns :: single sample supports all introns ERS025084, ERS025085 [ECO:0000348] ##Evidence-Data-END## COMPLETENESS: complete on the 3' end.



[View online »](#)

Synonyms: Di-Ras2


## Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DIRAS2 [RC200740], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DIRAS2 (1:2000). Positive lysates [LY413684] (100ug) and [LC413684] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffin-embedded Human adult brain tissue within the normal limits using anti-DIRAS2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human embryonic brain cortex tissue within the normal limits using anti-DIRAS2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffin-embedded Human embryonic cerebellum within the normal limits using anti-DIRAS2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.