

Product datasheet for TA809502S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TRMT12 Mouse Monoclonal Antibody [Clone ID: OTI3C10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C10 IHC. WB **Applications:**

Recommended Dilution: WB 1:2000, IHC 1:1000

Reactivity: Human Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human TRMT12 (NP_060426) produced in E.coli.

PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. Formulation:

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 50.1 kDa

Gene Name: tRNA methyltransferase 12 homolog

Database Link: NP 060426

Entrez Gene 55039 Human

Q53H54

Background: Wybutosine (yW) is a hypermodified guanosine at the 3-prime position adjacent to the

> anticodon of phenylalanine tRNA that stabilizes codon-anticodon interactions during decoding on the ribosome. TRMT12 is the human homolog of a yeast gene essential for yW

synthesis (Noma and Suzuki, 2006). [supplied by OMIM, Mar 2008]. COMPLETENESS:


complete on the 3' end.

TRM12; TYW2 Synonyms:






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TRMT12 ([RC203982], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TRMT12 (1:2000). Positive lysates [LY402634] (100ug) and [LC402634] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Ovary tissue within the normal limits using anti-TRMT12 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809502]) (1:1000)