

## **Product datasheet for TA809512**

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **HEF1 (NEDD9) Mouse Monoclonal Antibody [Clone ID: OTI6E3]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI6E3
Applications: WB

- 1 1-11 1 11-12

Recommended Dilution: WB 1:2000

**Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Full length human recombinant protein of human NEDD9 (NP\_006394) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 92.7 kDa

**Gene Name:** neural precursor cell expressed, developmentally down-regulated 9

Database Link: NP 006394

Entrez Gene 18003 MouseEntrez Gene 291044 RatEntrez Gene 4739 Human

Q14511






#### Background:

The protein encoded by this gene is a member of the CRK-associated substrates family. Members of this family are adhesion docking molecules that mediate protein-protein interactions for signal transduction pathways. This protein is a focal adhesion protein that acts as a scaffold to regulate signaling complexes important in cell attachment, migration and invasion as well as apoptosis and the cell cycle. This protein has also been reported to have a role in cancer metastasis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Aug 2012]

Synonyms: CAS-L; CAS2; CASL; CASS2; HEF1

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NEDD9 ([RC207200], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NEDD9 (1:2000). Positive lysates [LY401924] (100ug) and [LC401924] (20ug) can be purchased separately from OriGene.