

Product datasheet for TA809592

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ROR gamma (RORC) Mouse Monoclonal Antibody [Clone ID: OTI1G3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1G3

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-266 of human RORC

(NP_005051) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: RAR related orphan receptor C

Database Link: NP 005051

Entrez Gene 19885 MouseEntrez Gene 368158 RatEntrez Gene 6097 Human

P51449

Background: The protein encoded by this gene is a DNA-binding transcription factor and is a member of

the NR1 subfamily of nuclear hormone receptors. The specific functions of this protein are not known; however, studies of a similar gene in mice have shown that this gene may be essential for lymphoid organogenesis and may play an important regulatory role in

thymopoiesis. In addition, studies in mice suggest that the protein encoded by this gene may

inhibit the expression of Fas ligand and IL2. Two transcript variants encoding different

isoforms have been found for this gene. [provided by RefSeq, Jul 2008]






Synonyms: NR1F3; RORG; RZR-GAMMA; RZRG; TOR

Protein Families: Druggable Genome, Transcription Factors

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RORC ([RC206541], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RORC (1:2000). Positive lysates [LY401568] (100ug) and [LC401568] (20ug) can be purchased separately from OriGene.