

# Product datasheet for TA809611BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### **COL5A2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2H3]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2H3

**Applications:** WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 1200-1499 of human

COL5A2(NP\_000384) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 144.7 kDa

**Gene Name:** collagen type V alpha 2 chain

Database Link: NP 000384

Entrez Gene 1290 Human

P05997



### COL5A2 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2H3] - TA809611BM

Background:


This gene encodes an alpha chain for one of the low abundance fibrillar collagens. Fibrillar collagen molecules are trimers that can be composed of one or more types of alpha chains. Type V collagen is found in tissues containing type I collagen and appears to regulate the assembly of heterotypic fibers composed of both type I and type V collagen. This gene product is closely related to type XI collagen and it is possible that the collagen chains of types V and XI constitute a single collagen type with tissue-specific chain combinations. Mutations in this gene are associated with Ehlers-Danlos syndrome, types I and II. [provided by RefSeq, Jul 2008]

Synonyms: MGC105115

**Protein Families:** Druggable Genome

**Protein Pathways:** ECM-receptor interaction, Focal adhesion

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY COL5A2 (Cat# [RC213335], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-COL5A2 (Cat# [TA809611])(1:2000). Positive lysates [LY400138] (100ug) and [LC400138] (20ug) can be purchased separately from OriGene.