

Product datasheet for TA809616AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PASD5 (NPAS1) Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C9
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:500 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 319-522 of human

NPAS1 (NP_002508) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 62.5 kDa

Gene Name: neuronal PAS domain protein 1

Database Link: NP 002508

Entrez Gene 4861 Human

Q99742

Background: The protein encoded by this gene is a member of the basic helix-loop-helix (bHLH)-PAS family

of transcription factors. Studies of a related mouse gene suggest that it functions in neurons. The exact function of this gene is unclear, but it may play protective or modulatory roles during late embryogenesis and postnatal development. [provided by RefSeq, Jul 2008]

Synonyms: bHLHe11; MOP5; PASD5







Protein Families:

Druggable Genome, Transcription Factors

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NPAS1 ([RC207926], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NPAS1 (1:2000). Positive lysates [LY419279] (100ug) and [LC419279] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human embryonic brain cortex tissue within the normal limits using anti-NPAS1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809616]) (1:500)