

### **Product datasheet for TA809723**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **HDAC8 Mouse Monoclonal Antibody [Clone ID: OTI3F6]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI3F6
Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-243 of human HDAC8

(NP\_060956) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** histone deacetylase 8

Database Link: NP 060956

Entrez Gene 70315 MouseEntrez Gene 363481 RatEntrez Gene 55869 Human

Q9BY41

Background: Histones play a critical role in transcriptional regulation, cell cycle progression, and

developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes the deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large multiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isoforms have

been found for this gene. [provided by RefSeq, Oct 2009]






Synonyms: CDA07; CDLS5; HD8; HDACL1; MRXS6; RPD3; WTS

**Protein Families:** Druggable Genome, Transcription Factors

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY HDAC8 ([RC208390], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HDAC8 (1:2000) (TA809723). Positive lysates [LY413013] (100ug) and [LC413013] (20ug) can be purchased separately from OriGene.