

Product datasheet for **TA809809**

PD-L1 (CD274) Mouse Monoclonal Antibody [Clone ID: OTI13G7]


Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI13G7
Applications:	ELISA, FC, IF
Recommended Dilution:	IF 1:5000, FLOW 1:100
Reactivity:	Human (Does not react with: Mouse)
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 19-239 of human PD-L1/CD274(NP_054862) produced in HEK293 cells.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	31 kDa
Gene Name:	CD274 molecule
Database Link:	NP_054862 Entrez Gene 60533 Mouse Entrez Gene 29126 Human Q9NZQ7
Synonyms:	B7-H; B7H1; PD-L1; PDCD1L1; PDCD1LG1; PDL1
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Cell adhesion molecules (CAMs)




[View online »](#)


Product images:





PD-L1 ELISA with 13G7 Capture (TA809809) and 9E12 Detection ([TA808771]) Antibodies. Substrate used: Recombinant Human PD-L1 ([TP700201])





Immunocytochemistry staining of CD274 stable expression cells using anti-CD274 mouse monoclonal antibody ([TA507087]) (Left). The right is negative control (1:5000).





Flow cytometric Analysis of HCC78 cells, using anti-PDL1 antibody (TA809809), (Red), compared to isotype control, (green), and negative control (PBS), (Blue) (1:100).




Flow cytometric Analysis of stable expression CD274 cells using anti-CD274 antibody (TA809809) (Red) compared to a nonspecific negative control antibody (Blue). The left is 293T as negative control (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right)/untreated (Left) using anti-PDL1 antibody (TA809809) (1:100).



Flow cytometric analysis of living PBMCs treated with 10ug/ml PHA for 72h (Right) using anti-PDL1 antibody (TA809809). Cells incubated with a non-specific antibody (Left) were used as isotype control (1:100).



Detection of PDL1 neutralizing antibody using MACS column. GFP+/PDL1+ 293T cells (co-transfected with PDL1 and GFP plasmid [RC213071], PS10010) were incubated with either PDL1 antibody TA809809 (red), non-specific antibody (green), isotype control (blue) or PBS (black) and then mixed with PD1+ 293T cells [RC210364] linked with magnetic-beads. The mixed cells were pulled down using MACS column (Miltenyi Biotec) and analysed by Flow Cytometry. GFP+/PDL1+ cells would not be collected if PD1/PDL1 interaction is neutralized by the tested antibody (1:50).