

Product datasheet for TA809864AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CLCA1 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI8E7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8E7
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 344-412 of human

CLCA1(NP_001276) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: chloride channel accessory 1

Database Link: NP 001276

Entrez Gene 1179 Human

A8K7I4





Background:


This gene encodes a member of the calcium sensitive chloride conductance protein family. To date, all members of this gene family map to the same region on chromosome 1p31-p22 and share a high degree of homology in size, sequence, and predicted structure, but differ significantly in their tissue distributions. The encoded protein is expressed as a precursor protein that is processed into two cell-surface-associated subunits, although the site at which the precursor is cleaved has not been precisely determined. The encoded protein may be involved in mediating calcium-activated chloride conductance in the intestine. [provided by RefSeq, Jul 2008]

Synonyms: CACC; CaCC-1; CACC1; CLCRG1; GOB5; hCaCC-1; hCLCA1


Protein Families: Druggable Genome, Ion Channels: Other

Protein Pathways: Olfactory transduction

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CLCA1 (Cat# [RC218583], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CLCA1 (Cat# [TA809864])(1:2000).



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-CLCA1 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809864]) (1:500)