

Product datasheet for TA809878S

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

METTL7A Mouse Monoclonal Antibody [Clone ID: OTI4B10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4B10
Applications: IHC, WB

Recommended Dilution: WB 1:500~2000, IHC 1:2000

Reactivity: Human, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 34-104 of human

METTL7A(NP_054752) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28.1 kDa

Gene Name: methyltransferase like 7A

Database Link: NP 054752

Entrez Gene 315306 RatEntrez Gene 25840 Human

Q9H8H3


Background: Self-associates. [UniProtKB/Swiss-Prot Function]

Synonyms: AAM-B


Protein Families: Druggable Genome, Transmembrane



Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY METTL7A (Cat# [RC202601], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-METTL7A (Cat# [TA809878])(1:2000). Positive lysates [LY415471] (100ug) and [LC415471] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from 3 different cell lines by using anti-METTL7A monoclonal antibody (1:500).





Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-METTL7A mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809878]) (1:2000)



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-METTL7A mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809878]) (1:2000)