

## **Product datasheet for TA809899S**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com

OriGene Technologies, Inc.

EU: info-de@origene.com CN: techsupport@origene.cn

# **USP40 Mouse Monoclonal Antibody [Clone ID: OTI7D1]**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7D1
Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 767-912 of human

USP40(NP\_060688) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 141.4 kDa

**Gene Name:** ubiquitin specific peptidase 40

Database Link: NP 060688

Entrez Gene 55230 Human

O9NVE5





**Background:** Modification of cellular proteins by ubiquitin is an essential regulatory mechanism controlled

by the coordinated action of multiple ubiquitin-conjugating and deubiquitinating enzymes. USP40 belongs to a large family of cysteine proteases that function as deubiquitinating enzymes (Quesada et al., 2004 [PubMed 14715245]). [supplied by OMIM, Mar 2008].


##Evidence-Data-START## Transcript exon combination :: AJ583821.2 [ECO:0000332] RNAseq introns :: mixed/partial sample support ERS025081, ERS025082 [ECO:0000350] ##Evidence-

Data-END## COMPLETENESS: complete on the 3' end.

Synonyms: FLJ10785; FLJ42100

**Protein Families:** Protease

### **Product images:**



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-USP40 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA809899]) (1:150)