

Product datasheet for **TA809982**

N myc interactor (NMI) Mouse Monoclonal Antibody [Clone ID: OTI2C1]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI2C1
Applications:	WB
Recommended Dilution:	WB 1:500~2000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-223 of human NMI(NP_004679) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	34.9 kDa
Gene Name:	N-myc and STAT interactor
Database Link:	NP_004679 Entrez Gene 9111 Human Q13287
Background:	NMYC interactor (NMI) encodes a protein that interacts with NMYC and CMYC (two members of the oncogene Myc family), and other transcription factors containing a Zip, HLH, or HLH-Zip motif. The NMI protein also interacts with all STATs except STAT2 and augments STAT-mediated transcription in response to cytokines IL2 and IFN-gamma. The NMI mRNA has low expression levels in all human fetal and adult tissues tested except brain and has high expression in cancer cell line-myeloid leukemias. [provided by RefSeq, Jul 2008]




[View online »](#)


Synonyms: N-myc (and STAT) interactor; N-myc-interactor; N-myc and STAT interactor; N-myc interactor

Protein Families: Druggable Genome, Transcription Factors


Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NMI (Cat# [RC201543], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NMI (Cat# TA809982)(1:2000). Positive lysates [LY417813] (100ug) and [LC417813] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 3 different cell lines by using anti-NMI monoclonal antibody (1:500).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type A549 cells and NMI-Knockout A549 cells (KO, Cat# [LC800044]) were separated by SDS-PAGE and immunoblotted with anti-NMI monoclonal antibody TA809982 (1:500). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.