

### Product datasheet for TA810144BM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## SLC39A5 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI2F2]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2F2
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 48-178 of human

SLC39A5 (NP\_775867) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 56.1 kDa

**Gene Name:** solute carrier family 39 member 5

Database Link: NP 775867

Entrez Gene 283375 Human

Q6ZMH5


**Synonyms:** LZT-Hs7; MYP24; ZIP5

**Protein Families:** Transmembrane






# **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY SLC39A5 (Cat# [RC206665], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SLC39A5 (Cat# [TA810144])(1:2000). Positive lysates [LY406498] (100ug) and [LC406498] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-SLC39A5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810144]) (1:500)





Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-SLC39A5 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA810144]) (1:500)